



REGISTRATION NUMBER

--	--	--	--	--	--	--	--	--	--

# SRINIX COLLEGE OF ENGINEERING

## 2<sup>nd</sup> INTERNAL EXAMINATION 2021-22

Sub – Math-III

Branch - All

Full marks- 100

Time – 2.30hrs

### 1. Answer any all questions (Part – A)

( 2 x 10 =20 )

- What is the standard Deviation of Random variable ?
- Explain the order of convergence of an fixed iteration process.
- Define type-I and type-II error in hypothesis testing.
- What is multistep method?
- What do you mean by one-tail and two-tail testing?
- What is Newton's backward interpolation formula ?
- What is the idea of maximum likelihood method in estimating a parameter ?
- State the Baye's theorem.
- Distinguish between binomial and normal distribution.
- Explain the gauss quadrature formula.

### 2. Answer any eight questions (Part – B )

(6x8=48)

- Solve by Doolittle's method the system of equation
$$x_1 + x_2 + x_3 = 5$$
$$x_1 + 2x_2 + 2x_3 = 6$$
$$x_1 + 2x_2 + 3x_3 = 8$$
- Solve Numerically  $dy/dx = y-x$ , where  $y(0) = 2$ ;  $h = 0.1$ ; Find  $y(0.1)$  by Runge – kutta method of order 4.
- Solve by cholesky's method the system of equation
$$4x_1 + 10x_2 + 8x_3 = 44$$
$$10x_1 + 26x_2 + 26x_3 = 128$$
$$8x_1 + 26x_2 + 3x_3 = 214$$
- Suppose a large high school has 1100 female students and 900 male students. A random sample of 10 students is drawn with outreplacement. What is the probability exactly 7 of the selected students are female ?
- Bag A contains 3 red and 4 green balls. Bag B contains 4 red and 5 green balls. One ball is drawn at random from one of the bags and found to be red. What is the probability that it was drawn from bag A ?

f) Fit a straight line  $y=a+bx$  to the following data by the method of least square;

x	1	2	3	4	5
y	14	27	40	55	68

g)Using the Newton's divided difference formula calculate the value of  $f(2)$  from the following data;

x	1	3	4	6
F(x)	-3	9	30	132

h)Describe the Lagrange interpolation technique and find the value of  $f(10)$  for the given data.

x	5	6	9	11
F(x)	12	13	14	16

I)Find the real root of the equation  $\cos x - xe^x = 0$  correct up to three decimal place by using Newton Raphson method.

**3. Answer any two questions (Part – C)**

**(16 x 2 = 32)**

a) Find the correlation coefficient and the equation of the lines of regression for the following values of x and y

x	1	2	3	4	5
y	2	5	3	8	7

b) Solve the following system of equations by using Gauss-jacobi method

$$10x+y+z=12$$

$$x+10y+z=12, x+y+10z=12$$

c)Determine y for  $x=0.1,0.2,0.3,0.4,0.5$  where y is the solution of the differential equation  $dy/dx = 2(y+1); y(0)=0$  by using euler's method with  $h=0.1$ . find the exact solution and compare your numerical result with the exact solution.